



In Vitro Evaluation of NIH/3T3 Fibroblast Cell Viability After Treatment with Reundeu (*Staurogyne Elongata*) Leaf Extract

Fifi Hafizhatul Nuraini¹, Indah Puti Rahmayani Sabirin², Rhabiah El Fithriyah³, Euis Reni Yuslianti⁴

¹Faculty of Dentistry, Universitas Jenderal Achmad Yani, Indonesia

²Department of Oral Biology, Faculty of Dentistry, Universitas Jenderal Achmad Yani, Indonesia

³Department of Pedodontics, Faculty of Dentistry, Universitas Jenderal Achmad Yani, Indonesia

⁴Department of Oral Biology, Faculty of Dentistry, Universitas Jenderal Achmad Yani, Indonesia

Correspondence email: fifihafizhatull@gmail.com

ARTICLE INFO

Keywords:

Cytotoxicity, extract, fibroblast, Staurogyne elongata, WST-8 assay

History:

Received : 06/04/2026

Revision : 22/08/2026

Accepted : 23/08/2026

Published : 31/08/2026

Copyright © 2026 IJD

ISSN: 2775-0159



Open access under

[CC BY-SA 4.0](https://creativecommons.org/licenses/by-sa/4.0/)

International license.

ABSTRACT

Background: Reundeu (*Staurogyne elongata*) is a plant traditionally used as food and medicine, especially in the Java region. It contains bioactive compounds that have the ability to improve health. Therefore, it may show potential for formulation into standardized herbal medicine.

Objective: The aim of this study was to determine whether an ethanolic extract of reundeu (*Staurogyne elongata*) leaves exhibits a cytotoxic effect on NIH/3T3 fibroblast cells using a WST-8 assay.

Method: In this in vitro laboratory examination, the NIH/3T3 cell line, cultured in 96-well plates, was treated with reundeu leaves ethanolic extract at 3.125%, 1.56%, 0.78%, 0.39%, and 0.19% concentrations, with DMEM and a cell-control as controls. Cell viability was assessed using the WST-8 assay and measured at 490 nm using a spectrophotometer. Data were analyzed using one-way ANOVA to assess differences among groups, followed by Tukey's post-hoc test for multiple pairwise comparisons.

Outcome: The ethanolic extract at 3.125% concentration had mild toxicity to NIH/3T3 cells, but other concentrations showed high viability. One-way ANOVA demonstrated a significant effect of reundeu leaf extract concentration on NIH/3T3 cell viability ($p < 0.05$). Tukey's post-hoc analysis showed significant differences ($p < 0.05$) between the 3.13%, 1.56%, and 0.19% groups and other concentration groups.

Conclusion: The study showed that reundeu leaf ethanolic extract can be considered safe but mildly cytotoxic to fibroblasts at a concentration of 3.125%.

Citation (Vancouver style):

FH, Sabirin IPR, Fithriyah RE, Yuslianti ER. In vitro evaluation of NIH/3T3 fibroblast cell viability after treatment with Reundeu (*Staurogyne elongata*) leaf extract. Indones J Dent. 2026 Aug; 6(2), 158-169.

INTRODUCTION

The use of medicinal plants as primary constituents of therapeutic agents has been widely practiced in society since they are accessible, relatively low cost, and have an apparent safety profile. Plant-derived compounds continue to provide an important source of bioactive substances for medical or pharmaceutical development.^{1,2} To integrate herbal medicine into modern therapeutic treatment, it is necessary to conduct structured and continuous research to prove its effectiveness against disease, its level of safety for humans, and to recognize its cellular and molecular mechanisms.³ One of the plants that has been discovered for therapeutic purposes is reundeu (*Staurogyne elongata*).

The reundeu plant, also referred to as keji in certain areas of Central Java, has been traditionally used by society, especially in Java and Sumatra, as a food and medicinal plant. Empirically, the leaves and roots are either eaten directly or decocted in water for various health purposes such as joint pain remedy, kidney conditions, perinatal care, or canker sores.⁴⁻⁸ This traditional use highlights its potential as a candidate for advanced pharmacological and toxicological evaluation.

Phytochemical screening conducted on the leaves of this plant indicated the presence of secondary metabolites such as flavonoid compounds, saponins, tannins, polyphenols or phenols, and triterpenoids or steroids.^{7,8} The content of these phytochemicals can help protect the body from damage with antimicrobial, anti-inflammatory, and antioxidant effects.⁹⁻¹¹ A previous study by Noverita and Sinaga (2021) revealed that 50% reundeu leaf methanol extract has an inhibition zone against *Staphylococcus aureus* and *Bacillus cereus* with the agar diffusion method. Moreover, the active substance of reundeu leaves has strong antioxidant ability as shown by Marsiati and Patimah (2023), which indicated that the IC₅₀ of reundeu leaf water extract based on the DPPH test was 17.73 ppm.^{12,13}

Based on the observation of empirical uses and previous studies, Reundeu (*Staurogyne elongata*) leaf has considerable potential for development as medicinal agents; however, substances planned for therapeutic or dietary use must be evaluated for potential toxic or harmful effects. Cytotoxicity testing on cultured cells is commonly performed to assess the safety of bioactive materials prior to further development.^{14,15} These assays evaluate cellular responses, including changes in viability, proliferation, and cell damage following exposure to the tested substance.¹⁶

NIH/3T3 fibroblast cells are frequently used as an in vitro model to evaluate cytotoxic effects on normal cells, as they play a crucial role in tissue maintenance and wound healing and exhibit relatively stable growth characteristics. Therefore, assessment of fibroblast cell

viability provides valuable information regarding the biocompatibility of plant-derived extracts.¹⁷ In this study, cytotoxicity testing was conducted to evaluate the effect of reundeu leaf extract at various concentrations on fibroblast cell viability, aiming to determine a concentration range that is non-cytotoxic and potentially safe for further application.

Cell viability assays such as the WST (Water Soluble Tetrazolium)-8 assay are a common test employed to evaluate the cytotoxic effects of bioactive compounds *in vitro*. The WST-8 assay is a colorimetric method used to assess cell viability and examine whether fibroblast cell proliferation is affected by treatment.¹⁸ This study aims to examine the cytotoxicity of reundeu leaf ethanolic extract by measuring cell viability as an indicator of treatment-induced cell death *in vitro* on fibroblast cells.

RESEARCH METHODS

Leaf extract

Reundeu leaves were obtained from a reundeu cultivation area in Susuru, Ciamis, West Java, Indonesia. Plant identification was conducted at the Plant Taxonomy Laboratory of the Mathematics and Science Faculty, Padjadjaran University. Ethanol extracts of the leaves were prepared by maceration using 70% ethanol.¹⁹ Phytochemical compounds present in the extract were identified using specific reagents for each compound. The ethanolic extract of the leaves was subsequently formulated by dilution with distilled water to obtain concentrations of 3.13%, 1.56%, 0.78%, 0.39%, and 0.19%.

WST-8 assay materials

The 3T3 fibroblast cell culture used was the NIH 3T3 cell line cultured at the Aretha Cell Culture Laboratory, Bandung, from ATCC CL 183 P9 mouse embryonic cells. Fibroblast cells were seeded at a density of 5000 cells/well. Materials used in the assay were Dulbecco's Modified Eagle's Medium/DMEM (Elabscience) as a culture growth medium; WST-8 reagent (Elabscience); antimetabolic antibiotics; nanomycopulitin (Biowest); gentamycin; amphotericin; 0.25% trypsin EDTA (Merck); DMSO (Merck); PBS; a Biological Safety Cabinet (Thermo Scientific 1300 Series); a CO₂ incubator (Thermo Scientific); a 96-well plate (Costar); a microplate reader (Thermo Scientific); a micropipette (Finnpipette F2); an analytical balance (AXIS); a water bath; a refrigerator; a liquid nitrogen tank; and a microscope (Olympus CX31).

WST-8 assay

The prepared extracts will be tested using the WST-8 assay. Concentrations of 3.13%, 1.56%, 0.78%, 0.39%, and 0.19% on a 96-well plate were determined by purposive sampling.

This assay is based on the reduction of the WST-8 reagent, a water-soluble tetrazolium salt, by NADPH generated in viable cells. Formazan in WST-8 dissolved in tissue culture media. Subsequent quantitative spectrophotometry measurement was evaluated at a 490 nm wavelength, and the absorbance result will determine the viability of the cells. The cells in the well plate will visibly have an orange color.

The cytotoxicity assay was performed after preparation of the ethanolic extract of reundeu leaves at the previously explained concentrations as treatment. 3T3 fibroblast cells were cultured and prepared, thawed, and maintained under required conditions. Cells were seeded in 96-microplates at 5,000 cells per well, and the volume was 100 µL per well. Afterward, they were incubated at 37°C in a condition of 5% CO₂ until the confluence reached 80%. This assay was performed in triplicate.

Cells in the well plate were treated with serial concentrations of the extract, similarly for control treatments, then incubated for 24-48 hours, followed by the addition of WST-8 reagent and further incubation for 3-4 hours. Measurement of absorbance was done with an ELISA reader at a 490 nm wavelength, followed by the calculation of cell viability based on the absorbance values.

RESEARCH FINDINGS

Fibroblast cell viability was assessed using the WST-8 assay, with spectrophotometric measurements of absorbance obtained at a wavelength of 490 nm. Fibroblast cell cultures were exposed to reundeu (*Staurogyne elongata*) leaf extract at concentrations of 3.13%, 1.56%, 0.78%, 0.39%, and 0.19%. The observed cell viability values at 0.19%, 0.39%, 0.78%, and 1.56% concentrations remained above 90%, which is commonly indicated as non-cytotoxic or safe. In contrast, the lowest cell viability was observed at the highest concentration (3.13%); however, this reduction remained within the range of mild cytotoxicity, according to the International Organization for Standardization (ISO) 10993-5 standard, which divides toxicity levels based on the percentage of cell viability. It interprets 100%-80% viability as non-toxic, 60-80% as mild toxicity, 40-60% as moderate toxicity, and less than 40% as very toxic. The results of the WST-8 assay and cell viability percentages are presented in Table 1 and Figure 2, respectively.

Table 1. Viability of fibroblast cells with treatment from each group

Treatment	Viability (%)	Inhibition (%)	p-value
Negative control	100.00 ± 1.56	0.00 ± 1.56	0.000*
1% DMSO	98.42 ± 3.00	1.58 ± 3.00	
SE 3.125%	79.62 ± 2.07	20.38 ± 2.07	
SE 1.56%	90.72 ± 0.88	9.28 ± 0.88	
SE 0.78%	97.03 ± 1.32	2.97 ± 1.32	
SE 0.39%	100.57 ± 0.82	-0.57 ± 0.82	
SE 0.19%	106.13 ± 2.92	-6.13 ± 2.92	

Explanation:

* : ANOVA test shows $p < 0.05$ considered significant;

SE: *Staurogyne elongata* extract




Figure 1. Viability of 3T3 fibroblast cells after treatment with Negative control, DMSO, and *S. elongata* leaves (SE) at various concentration as calculated using the WST-8 assay. Data are presented as mean ± SD.

The results of the Shapiro-Wilk normality test and Levene's test showed that the data were normally distributed and homogeneous ($p > 0.05$). Statistical analysis using a One-Way ANOVA test showed a significant effect of reundeu leaf extract concentration on NIH/3T3 fibroblast cell line viability ($p < 0.0001$). The differences between treatment groups at various concentrations were further analyzed with a Post Hoc Tukey HSD test. The analysis showed that several treatment groups differed significantly, while no significant differences were observed among the negative control group, DMSO, and lower extract concentrations, as presented in Table 2. Furthermore, microscopic examination confirmed these findings, revealing that the viable cells retained a light or shiny appearance across the control and treatment groups, with no prominent morphological degradation observed at lower concentrations (Figure 1).

Table 2. NIH/3T3 fibroblast cell viability difference among all groups

	NC	1% DMSO	SE 3.125%	SE 1.56%	SE 0.78%	SE 0.39%	SE 0.19%
NC	-	0,911	0,000*	0,000*	0,377	1,000	0,004*
1% DMSO		-	0,000*	0,000*	0,951	0,721	0,000*
SE 3.125%			-	0,000*	0,000*	0,000*	0,000*
SE 1.56%				-	0,003*	0,000*	0,000*
SE 0.78%					-	0,200	0,000*
SE 0.39%						-	0,010*
SE 0.19%							-

Explanation:

* : Tukey HSD post-hoc test with $p < 0,05$ showed that the difference considered significant.

NC: negative control; SE: *Staurogyne elongata* extract.




Figure 2. NIH/3T3 fibroblast cells examination with light microscope of 100x magnification with description as follows: A. Negative control; B. DMSO group; C. Application of 3.125% of reundeu leaf extract; D. Group with application of 1.56% extract; E. Group with application of 0.78% extract; F. Group with application of 0.39% extract; G. Group with application of 0.19% extract. The viable cells showed a light or shiny appearance, while unalive cells would appear dark.

DISCUSSION

Reundeu (*Staurogyne elongata*) is a plant widely used as a medicinal plant in Indonesia. Based on phytochemical tests, its leaves contain various bioactive compounds, such as alkaloids, flavonoids, saponins, polyphenols, and steroids or triterpenoids. Although the phytochemical compounds of reundeu leaves have various therapeutic potentials, evaluating their cytotoxic effect is essential to support their safe usage as herbal medicine.⁷

Reundeu (*Staurogyne elongata*) is a plant widely used as a medicinal plant in Indonesia. Based on phytochemical tests, its leaves contain various bioactive compounds, such as alkaloids, flavonoids, saponins, polyphenols, and steroids or triterpenoids. Although the phytochemical compounds of reundeu leaves have various therapeutic potentials, evaluating their cytotoxic effect is essential to support their safe usage as herbal medicine.⁷

Based on the study results, reundeu leaves affected fibroblast cell viability in a concentration-dependent manner, as assessed using the WST-8 assay. The viability of fibroblast cells at the highest concentration (3.13%) exhibited mild cytotoxicity with reduced cell viability compared to other groups. In contrast, lower concentrations showed viability comparable to the negative control and DMSO solvent groups, indicating that lower doses had minimal cytotoxic effects. Statistical tests at concentrations of 0.78% and 0.39% did not differ significantly from the control cell group and the DMSO solvent group. These findings suggest that reundeu leaf extract is concentration-dependent and relatively safe for fibroblast cells at lower concentrations.

The cytotoxic effect observed at higher concentrations may be attributed to the increased levels of bioactive compounds. While these compounds present antioxidant properties at low doses, they can induce cellular stress and interfere with mitochondrial metabolic activity at higher concentrations. This is particularly relevant to the WST-8 assay, which reflects the activity of mitochondrial dehydrogenase as a parameter.^{19,20}

Reundeu leaves contain phytochemical constituents such as alkaloids, flavonoids, saponins, polyphenols, and steroids or triterpenoids. These have recognized health benefits due to their antioxidant, antibacterial, and anti-inflammatory properties, as well as their protective properties against free radicals.^{10,21-23} Several phytochemicals, including flavonoids, saponins, and alkaloids, have toxic potential and can contribute to reduced viability through mechanisms like inducing oxidative stress, cell membrane damage, and cell cycle inhibition.^{9,23-27} These compounds have the potential to cause adverse biological effects or toxicity reactions leading to cell death when applied in high concentrations. Therefore, it is important to comprehend the safe dosage if it is aimed to be developed in the field of treatment.²⁸⁻³⁰

The dose-dependent cytotoxicity observed in this study is coherent with findings from an acute oral toxicity study of reundeu leaves. This study reported no significant toxic effects in terms of changes in weight, autonomic system, behavior, sensory, neuromuscular, cardiovascular, respiratory, eye, gastrointestinal, urinary, and skin of mice, and no mice died during the study, indicating an LD50 value greater than 5 g/kg body weight. However, in vivo toxicity outcomes cannot be directly compared with in vitro cytotoxicity results. These findings suggest that differences in metabolic processing and exposure conditions may account for the mild cytotoxicity at higher concentrations in fibroblast cell cultures in this study.³¹

The relationship between the cytotoxicity of herbal extracts, their concentration, and their phytochemical content has been established through studies on cell viability after exposure to various concentrations of plant extracts. A study by Zain (2020) on the cytotoxicity of oil palm leaf on the fibroblast 3T3 cell line using an MTT assay showed that cell line viability at all concentrations was above 70% and also increased viability at lower concentrations.¹⁵ Tyatana (2021) demonstrated that the seed coat extract of *B. flabellifer* L. exhibited a concentration-dependent cytotoxic effect on fibroblasts with an MTT assay; concentrations from 1.56% to 12.5% were classified as moderately cytotoxic, while higher concentrations displayed severe toxicity.³² Based on the study by Laksmiawati (2016) regarding the effect of *G. vulgaris* and *A. muricata* extracts on the viability of macrophage cell line RAW264.7 with an MTS assay, higher concentrations of extracts were cytotoxic, while the application of lower extract concentrations indicated non-toxic levels and higher cell viability.³³ The results of these studies are in line with those of Gothai (2016), who also showed that the toxicity rate of a material is directly proportional to its concentration.³⁴ However, Harjadi et al. (2022) found that administration of tamarind ethanol extract on 3T3-L1 cells with a WST-8 assay at concentrations of 0.001, 0.004, 0.016, 0.063, 0.25, and 1.00 µg/mL exhibited low toxicity, with a viability level of 85%. These study results indicated that a minimal cytotoxic effect occurred at lower concentrations, but cell viability reduced with increased plant extract concentration. However, the limitations of this study were that the concentration range was less varied, the solvent used was only ethanol, and only a single method of viability assay was performed. In addition, in vitro cytotoxicity testing has another limitation since it cannot completely represent complex biological processes such as systemic effects, metabolism, and cell interactions in tissues that can be examined in vivo.^{20,34}

CONCLUSION

This research indicated that the ethanol extract of reundeu (*Staurogyne elongata*) leaf exhibits mild to minimal cytotoxicity at the tested concentrations. The implication of this study suggests the potential of reundeu leaves for further development as a safe natural ingredient. These findings may give preliminary support for the use of reundeu leaf extract for alternative therapeutic applications, such as in wound healing. Therefore, from these results, further investigations, such as in vivo testing and clinical evaluations, are required to assess the comprehensive level of safety and efficacy of reundeu leaf extract for therapeutic use.

DECLARATION

Authors Contributions

[IS]: Conceptualization, Methodology, Formal Analysis, Writing – Original Draft; [FN]: Investigation, Software, Data Curation; [EY]: Supervision, Writing – Review & Editing, [RF]: Writing – Review & Editing. All authors have read and approved the published version of the manuscript and agreed to be accountable for all aspects of the work.

Conflict of Interest

authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Funding Statements

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Ethics Approval and Consent to Participate

The study protocol was reviewed and approved by the Ethics Committee of Faculty of Medicine, Universitas Jenderal Achmad Yani (Protocol No. 006/UH.3/02/2025).

Consent for Publication

Not applicable.

Data Availability Statement

The data supporting the findings of this study are available from the corresponding author upon reasonable request.

Declaration of Generative AI In Scientific Writing

The authors declare that no generative AI or AI-assisted technologies were used in the writing or preparation of this manuscript.

Acknowledgments

Authors The authors would like to thank Faculty of Dentistry, Universitas Jenderal Achmad Yani for support, and the staff of Aretha Cell Culture Laboratory, Bandung, Indonesia, for analytical assistance.

REFERENCES

1. Moghadam, Moghadam ET, Yazdani M, Tahmasebi E, Tebyanian H, Ranjbar R, Yazdani A, et al. Current herbal medicine as an alternative treatment in dentistry: In vitro, in vivo and clinical studies. *Euro Jour Pharm. Elsevier B.V.* 2020; 889(173665). doi: 10.1016/j.ejphar.2020.173665 PubMed PMID: 33098834.
2. Tahir AH, Hussain Z, Yousuf H, Fazal F, Tahir MA, Kashif M. Traditional herbal medicine and its clinical relevance: a need to preserve the past for the future. *J Biosci Med (Irvine)*. 2022;10(07):64–75. doi:10.4236/jbm.2022.107005.
3. Etaware P, America O, Egara, O, Ekun V. Herbal medicine: Scientific validation and future prospects. *Int J Pharm and Chem*. 2025; 11(3):67-75. doi: 10.11648/j.ijpc.20251103.12.
4. Safitri H, Malik A, Mukaromah AS. Secondary metabolite profile in stem and root of keji plant (*Staurogyne elongata* (Blume) Kuntze). *Biotropika: Journal of Tropical Biology*. 2024; 12(1), 8–15. doi: 10.21776/ub.biotropika.2024.012.01.02.
5. Dewi AP, Peniwidiyanti P, Hariri MR, Hutabarat PWK, Martiansyah I, Lailaty IQ, et al. Ethnobotany of food, medicinal, construction and household utilities producing plants in Cikaniki, Gunung Halimun Salak National Park, Indonesia. *J Mount Sci*. 2023; 20(1): 163–181. doi:10.1007/s11629-021-7108-5
6. Handayani A. Utilization of medicinal plants by people around Gunung Simpang Nature Reserve, West Java. *Pros Sem Nas Masy Biodiv Indon*. 2015; 1(6):1425-1432. doi:10.13057/psnmbi/m010628.
7. Maulani MI, Purwanti L, Dasuki UA. Antibacterial activity test of reundeu (*Staurogyne elongata* (Bl.) O.Kuntze) leaves extract against *Staphylococcus aureus* and *Escherichia coli*. *Pros Farmasi Unisba*. 2017; 3(2):565-569.
8. Iskandar J. Pandemics and sustainable human nature relations: A case study in Baduy community, South Banten, Indonesia. *E3S Web of Conferences*. 2021; 249: 01001.
9. Mariani R. Isolasi senyawa turunan fenol dari herba reundeu (*Staurogyne elongata* (Blume) O.Kuntze). *Repository. Universitas Garut, Garut*. 2014.
10. Akhmadi C, Utami W, Annisaa' E. Narrative review: senyawa fitokimia dan aktivitas farmakologi family Basselaceae sebagai obat luka. *Journal of Research in Pharmacy*. 2022 Sep;2(2):77–85.
11. Cedillo-Cortezano M, Martinez-Cuevas LR, López JAM, Barrera-López IL, Escutia-Perez S, Petricevich VL. Use of medicinal plants in the process of wound healing: A literature review. *Pharm*. 2024; 17(303):1–27. doi:10.3390/ph17030303.
12. Noverita N, Sinaga E. Antibacterial bioactivity from extract of reundeu caret (*Staurogyne longata* [sic]) and honje (*Etlingera hemisphaerica*). *J Trop Biodiv*. 2021; 2(1): 21-32.
13. Marsiati H, Patimah WA. Antioxidants and aging: combination and single tea on reundeu leaves (*Staurogyne elongata* (Ness) Kuntze) and Ulam Raja (*Cosmos caudatus*). *J Int Aging Mov*. 2023;5(1):29-38.
14. Ghica A., Tănase M L, Niculițe CM, Tocilă A, Popescu L, Luță EA, et al. In vitro toxicity evaluation of some plant extracts and their potential application in xerosis cutis. *Cosmetics*. 2024; 11(4):1-22. doi:10.3390/cosmetics11040124
15. Zain MSC, Lee SY, Sarian MN, Fakurazi S, Shaari K. In vitro wound healing potential of flavonoid c-glycosides from oil palm (*Elaeis guineensis* jacq.) leaves on 3T3 fibroblast cells. *Antioxidants*. 2020; 9(4):1-18. doi:10.3390/antiox9040326
16. Soltanian S, Sheikhabaei M, Mohamadi N. Cytotoxicity evaluation of methanol extracts of some medicinal plants on P19 embryonal carcinoma cells. *J of App Pharm Sci*. 2017;7(7): 142–149. doi:10.7324/JAPS.2017.70722
17. Rahimi AM, Cai M, Hoyer-Fender S. Heterogeneity of the NIH3T3 fibroblast cell line. *Cells*. 2022; 11(17):1-19. doi: 10.3390/cells11172677
18. Buranasin P, Mizutani K, Iwasaki K, Pawaputanon N, Mahasarakham C, Kido D, et al. High glucose-induced oxidative stress impairs proliferation and migration of human gingival fibroblasts. *PLoS ONE*. 2018; 13(8): e0201855. doi: 10.1371/journal.pone.0201855
19. Maryam F, Utami YP, Mus S, Rohana R. Perbandingan beberapa metode ekstraksi ekstrak etanol daun sawo duren (*Chrysophyllum cainito* L.) terhadap kadar flavanoid total menggunakan metode

- spektrofotometri UV-VIS. Jurnal Mandala Pharmacon Indonesia. 2023; 9(1), 132–138. doi:10.35311/jmpi.v9i1.336,
20. Harjadi KR, Ishida M, Nishi K, Sugahara T, Putra ABN. Water and ethanol extracts of tamarind (*Tamarindus indica*) suppress lipid accumulation in 3T3-L1 cells. Canrea Journal: Food Technology, Nutritions, and Culinary Journal. 2022; 5(1):1-11. doi: 10.20956/canrea.v5i1.630
21. Fauziah N, Maulidiyah M, Hartanto TP, Putri SND, Sabhira AS, Mukarromah IW, et al. Artikel review: Studi fitokimia dan farmakologi tanaman kelor (*Moringa oleifera* Lam). The Journal General Health and Pharmaceutical Sciences Research. 2023;1(4): 45–52.
22. Pengelly A. Triterpenoids and saponins. In The constituents of medicinal plants. CABI. 2021:95-111. doi: 10.1079/9781789243079.0006
23. Dalimunthe A, Anjelisa P, Hasibuan Z, Silalahi J, Sinaga SF, Satria D. Antioxidant activity of alkaloid compounds from *Litsea cubeba* Lour. OJCHEG. 2018; 34(2):1149-1152. doi:10.13005/ojc/340270
24. Nugraha MFI, Novita H, Rajamuddin MA, Yunita R, Enggarini W, Maharani FH, et al. Analisa fitokimia dan aktivitas antibakteri *Staurogyne* sp. pada bakteri penyakit ikan / Phytochemical analysis and antibacterial activity of *Staurogyne* sp. in fish disease bacteria. Torani: Journal of Fisheries and Marine Science. 2019 Jun;2(2):77–86.
25. Bournine L, Bensalem S, Fatmi S, Bedjou F, Mathieu V, Mokrane Iguer-Ouada M, et al. Evaluation of the cytotoxic and cytostatic activities of alkaloid extracts from different parts of *Peganum harmala* L. (Zygophyllaceae). Eur J Integr Med. 2017 Jan; 9:91-996. doi: 10.1016/j.eujim.2016.10.002
26. Khan MI, Karima G, Khan MZ, Shin JH, Kim JD. Therapeutic effects of saponins for the prevention and treatment of cancer by ameliorating inflammation and angiogenesis and inducing antioxidant and apoptotic effects in human cells. In J Mol Sci. 2022; 23(18):1-15. doi:10.3390/ijms231810665
27. Hidayati DN, Safitri EI, Yumni GG, Salsabiila I, Rahayu P. Cytotoxic activity of *Eugenia polyantha* Wight young leaves purification extract and fraction on T47D. 2023;13(2):131–139. doi:10.22435/jki.v13i2.6607
28. Ghosh P, Das C, Biswas S, Nag SK, Dutta A, Biswas M, et al. Phytochemical composition analysis and evaluation of in vitro medicinal properties and cytotoxicity of five wild weeds: A comparative study. F1000Research. 2020; 9(493);1–23. doi:10.12688/f1000research.22966.1
29. Albino AM, Gomes DL, dos Santos AH, Malta GSL, de Oliveira DHL, Souto LFL, et al. Cytotoxicity evaluation and dereplication of flavonoids-guided by antioxidant activity and total phenolics content from *Ephedrantus amazonicus* leaves. Brazilian Archives of Biology and Technology. 2024;67(e24231132):1–10. doi:10.1590/1678-4324-2024231132
30. Takayama K, Ito K, Matsui A, Yamashita T, Kawakami K, Hirayama D, et al. In vivo gene expression profile of human intestinal epithelial cells: From the viewpoint of drug metabolism and pharmacokinetics. Drug Metab Dispos. 2021 Mar; 49(3):221-232. doi: 10.1124/dmd.120.000283
31. Hermanto F, Sutjiatmo AB, Vikasari SN, Faramayuda F, Syam AK., et al. Uji toksisitas akut ekstrak etanol herba reundeu (*Staurogyne elongata*) pada mencit galur Swiss Webster. Pros Per Il Ikatan Apoteker Indonesia. 2019; ISBN: 978-979-95108-9-1
32. Tyatana N, Sudiono R, Janti J. Cytotoxicity effect of *Borassus flabellifer* L. seed coat on fibroblast. Journal of Indonesian Dental Association. 2021; 4(2):83–89. doi:10.32793/jida.v4i2.681
33. Laksmiawati DR, Prasanti AP, Larasinta N, Syauta GA, Hilda R, Ramadaniati HU, et al. Anti-inflammatory potential of gandarusa (*Gendarussa vulgaris* nees) and soursoup (*Annona muricata* L) extracts in LPS stimulated-macrophage cell (RAW264.7). Journal of Natural Remedies. 2016; 16(2): 73–81. doi:10.18311/jnr/2016/5367
34. Gothai S, Arulselvan P, Tan WS, Fakurazi S. Wound healing properties of ethyl acetate fraction of *Moringa oleifera* in normal human dermal fibroblasts. J of Intercultural Ethnopharm, 2016; 5(1): 1–6. doi:10.5455/jice.20160201055629